

Determining the β -function of the strong interaction and closing the light gluino window

F. Csikor⁽¹⁾ and Z. Fodor^{(2)*}

⁽¹⁾ *Institute for Theoretical Physics, Eötvös University, H-1088 Budapest, Hungary*

⁽²⁾ *Theory Division, CERN, CH-1211 Geneva 23, Switzerland*

We discuss the perturbative running of α_s in a model independent way. Our analysis contains data on the hadronic τ decay and hadronic cross sections in e^+e^- annihilation between 5 GeV and M_Z . We determine the color coefficients and the perturbative β -function of the strong interaction. The results are in agreement with QCD and rule out the QCD+light gluino scenarios on the 70.8 – 93.0% CL. We combine our method with that of the multi-jet analysis at LEP. The combined result rules out light gluinos at least on the 99.76% CL, provided that nonperturbative effects are not large.

Asymptotic freedom is one of the most interesting predictions of QCD. In order to study the running of α_s , one can collect its values at different scales (e.g. τ decay, DIS, Υ decay, e^+e^-) and compare them [1]. Another way is to analyse a single experiment (e.g. $\bar{p}p$, ep or τ decay [2]). Nice agreement has been found between experiments and QCD. Some differences between the results obtained from low-energy and high-energy experiments led to a number of speculations; e.g. that the apparent slower running of α_s could be due to additional light fermions of the theory.

Supersymmetric phenomenology deals normally with sparticles of masses $\mathcal{O}(100\text{GeV})$. The only exception is the light gluino [3] with mass ≤ 1.5 GeV (window I) [4,5] and 3 – 5 GeV (window III) [6]. We will discuss window I and window III separately. Since such a light particle influences the running of α_s the comparison of high and low energy experiments could open or close the light gluino window [7]–[9]. A consistent analysis must contain the virtual gluino effects not only in the running (technically in the β -function), but in all loop diagrams, extracting α_s from the experimental results, too [9].

One can even determine the β -function of the strong interaction as done in ref. [2]. In order to extract the three-loop coefficients one needed large values of α_s , that is the small energies of τ decay. The result is in good agreement with QCD.

It has been suggested [10] to look for gluinos in four-jet events. The LEP collaborations determined the color coefficients (C_A/C_F and T_F/C_F) in multi-jet events. The results (c.f. [1]) are in good agreement with QCD. An extension of QCD including light gluinos would result in a somewhat different T_F/C_F . The four-jet cross-section is known only at tree-level. Without the knowledge of the loop corrections this method alone can hardly give a clear answer to the question of light gluinos.

In this letter we determine the color coefficients and the perturbative β -function of the strong interaction from experimental results. We consider gauge theories with fermions in the fundamental representation and their extended versions with light gluinos. We perform an analysis on the three-loop level in the $\overline{MS}$ scheme. The experimental inputs are R_τ and R_h for energies between 5 GeV and M_Z . We combine our method with that of the multi-jet analysis at LEP.

One of the clearest ways to study the running of the coupling, thus the β -function, of the theory would be to study one given experimental quantity at different energies. We chose R_h 's at energies larger than 5 GeV, which are very clear quantities both experimentally and theoretically, due to their minimal nonperturbative corrections. Moreover, these are the only quantities known to $\mathcal{O}(\alpha_s^3)$ relative to the leading order, thus only a minimal scale ambiguity is present. In order to study the running of α_s one needs precise R_h measurements for a large energy region. Unfortunately, R_h measurements are limited by their small statistics for energy scales below M_Z . For this reason we have also considered another, strongly related (still not very low scale) quantity R_τ in our analysis. Since we are aware of the theoretical criticism on α_s determinations based on R_τ we include the effects of all known uncertainties in case of QCD [11]–[13]. The nonperturbative estimates to R_τ in the light gluino case might be unreliable. Therefore, we do not include R_τ in our analysis for window I gluinos. (Window III gluinos are much too heavy to contribute to R_τ .)

Let us suppose that the strong interaction is described by a gauge theory based on a simple Lie-group, which fixes the color coefficients of the theory (C_F, C_A, T_F). The three basic processes in the theory are: gluon bremsstrahlung from a quark, splitting of a gluon into two gluons and splitting of a gluon into two quarks. To lowest order their

amplitudes are proportional to a universal coupling and to C_F , C_A and T_F , respectively. E.g. for an $SU(N)$ gauge group one has $C_F = (N^2 - 1)/(2N)$, $C_A = N$ and $T_F = 1/2$. The β -function of the strong interaction, hadronic cross-sections and widths are calculated in terms of these coefficients and the active number of fermions. We will look for a set of parameters which describes the experimental data most accurately and give the corresponding confidence level (CL) regions. The outcome –that is the best fit– does not necessarily predict a meaningful theory. Nevertheless, it could tell the difference between QCD and QCD+light gluino scenario.

The β -function: $da(\mu)/d\log\mu = -\beta_0 a^2 - \beta_1 a^3 - \beta_2 a^4$ is known upto three-loop order. We factorize out the inconvenient π -s by using $a = C_F \alpha_s/(2\pi)$. Introducing $x = C_A/C_F$ and $y = T_F/C_F$ gives $\beta_0 = 11x/3 - 4(n_f y + x n_{\tilde{g}}/2)/3$, where n_f is the number of active flavours and $n_{\tilde{g}}$ is the number of active gluinos. The expressions for β_1 and β_2 are more complicated [14,15]. In the running of α_s we follow ref. [16] for threshold-effects. We solve the above renormalization group equations exactly.

The other important quantity in the analysis –the hadronic cross section in e^+e^- annihilation via a virtual photon– is known on the three-loop level, too.

$$\begin{aligned} R_\gamma &= \sigma(e^+e^- \rightarrow \gamma \rightarrow \text{had})/\sigma_o(e^+e^- \rightarrow \gamma \rightarrow \mu\mu) \\ &= 3\Sigma_f q_f^2(1 + K_1 a + K_2 a^2 + K_3 a^3 + \dots), \end{aligned} \quad (1)$$

where $K_1 = 3/2$. K_2 , K_3 is more complicated and also known for arbitrary C_F , C_A and T_F [14,17]. The hadronic decay ratio of the τ is defined: $R_\tau = \Gamma(\tau^- \rightarrow \nu_\tau + \text{had})/\Gamma(\tau^- \rightarrow \nu_\tau e^- \bar{\nu}_e)$. Its perturbative value is strongly related to R_γ . For a recent analysis and a review on R_τ see eg. [12] and [18]. The hadronic decay width of the Z boson $R_Z = \Gamma(Z \rightarrow \text{had})/\Gamma(Z \rightarrow \mu^+\mu^-)$ is again a function of R_γ and known at the three-loop level. At the energy scales of the analysis important electroweak and mixed corrections appear. We include these corrections.

The experimental values for R_τ are given in [19] with an average of $R_\tau = 3.616 \pm .02$. Averaging for the four LEP experiments and three leptons [20] gives $R_Z = 20.778 \pm .029$. The references for hadronic cross sections at energies below M_Z are taken from [21]. We have collected all the existing published data, and some unpublished results, too. Some of them were binned by the experimental groups. The total number of data points included in our analysis is 182.

Combining results from different experiments is a delicate question, with major problems. i) The use of the radiative corrections is not unique, reflecting the state of art at the time of publication. ii) The results do depend on the mass of the Z boson and top-quark –which was assumed to be approximately 20 GeV at the early eighties. iii) The measurement of the total cross section was not performed in the full phase space, and Monte-Carlo acceptance calculations based on different assumptions were done. We corrected for the first two problems, correction for the third one is practically impossible. For further details see [22]– [23].

There are different sources of uncertainties in the determination of α_s . We treat them in a unified manner. We add the systematic errors linearly. The total systematic error estimates and the statistical errors are combined quadratically. The overall normalization errors within one experiment are correlated. We include these correlations. We minimize $\chi^2 = \Delta^T V^{-1} \Delta$, where Δ is an n -vector of the residuals of $R_i - R_{fit}$ for the n individual results and V is an $n \times n$ error matrix. In V , the diagonal elements V_{ii} are the squares of the total errors for the i th measurement and the off diagonal elements V_{ij} correspond to the correlations between the i th and j th measurements. For different points from the same experiment V_{ij} is given by the product of the normalization errors [22]. The separation of the systematics into point-to-point and overall normalization error is ambiguous. Our results are rather insensitive to a moderate change of this splitting in a given experiment. For some experiments the above separation was not even explicitly given. We checked that our results are stable against even large variations of the splitting in these cases. However, the unreasonable extreme case of totally uncorrelated fitting would give rather different results with much more predictive power. We have assumed that the results of different experiments are uncorrelated. In order to perform a further consistency check we included an additional hypothetical overall error of 1% in the correlation matrix. The change of the result turned out to be negligible. Since the experimental groups usually give their results with binning in fixed energy intervals we checked that further binning had practically no influence on our results.




FIG. 1. The 68.3% CL regions for $x = C_A/C_F$ and $y = T_F/C_F$. The QCD value is represented by a square.

Due to the relatively poor statistics for the e^+e^- experiments below M_Z the theoretical uncertainties are dominant only for R_τ and non-negligible for R_Z . Estimating the nonperturbative and perturbative errors in τ -decay we use the results of [11,12]. For a fixed order QCD calculation we assume that the error is equal to the last computed term. Including all the errors we get $R_\tau = 3.616 \pm .143$, which corresponds to $\alpha_s(M_\tau) = .335 \pm .053$ and $R_Z = 20.778 \pm .0387$, which corresponds to $\alpha_s(M_Z) = .123 \pm .006$. We assume that currently incalculable and/or model dependent nonperturbative corrections are negligible or correctly estimated. If the nonperturbative corrections turn out to be larger than our estimates, the confidence levels for light gluino exclusion to be determined below should be lowered. Note that both error estimates are certainly very conservative.

Having included all the errors one can determine the best fit values and the CL regions for $x = C_A/C_F$ and $y = T_F/C_F$ with or without light gluinos.

Fig. 1 contains the 68.3% CL regions for theories without gluinos and with window III gluinos. As it can be seen the experimental results are in agreement with QCD; however, they rule out the QCD+light gluino scenario on the 93 (90.7)% CL's for $M_{\tilde{g}} = 3$ (5) GeV, respectively. The variation of the fixed χ^2 boundaries is quite small for the above mass region. For binning level 50 (d.o.f. 48) the χ^2 values are 37.37 without and 37.94 with gluinos ($M_{\tilde{g}} = 3$ GeV) for the best fits. As mentioned before, for window I gluinos we do not include R_τ in our study. In this case we have a 70.8% exclusion only.




FIG. 2. The 68.3% and 95.4% CL regions for β_0 and β_1 .

Note the important difference between the presentations of our result and that of the 4-jet analysis for the case with light gluinos [1]. In our case the best fits and the CL contours for the x, y variables are determined separately for the theories with and without light gluinos. These results are compared for both theories with the QCD values: $\bar{x} = 2.25$, $\bar{y} = .375$. Due to the simplicity of the tree-level treatment of the 4-jet analysis in ref. [1], it was possible to produce the same best fits and contours for both theories and compare this unique result with $\bar{x}, \bar{y}$ above in the theory without light gluinos, and with $\bar{x}$ and with an effective $\bar{y}_{eff} = .6$ for the gluino extended theory. The 4-jet results can easily be expressed in terms of our variables.

Two remarks are in order: i) the 4-jet analysis is based on tree-level calculations, whereas our method contains corrections up to three-loops. (In [1] also the one-loop analysis to 2- and 3-jets is done; however, the errors on x, y are huge.) ii) our method contains the running of α_s , too, thus it gives a simultaneous check of the β -function.

Assuming [24] that the underlying gauge group is SU(3), we may consider the number of gluinos as a free parameter. The best fits give $n_{\tilde{g}} = -.638 \pm 1.17$ for window I and $n_{\tilde{g}} = .0078 \pm .52$ for window III. The CL's are given by Bayes' theorem allowing only non-negative integer $n_{\tilde{g}}$: 72.5% for window I and 87.7% for window III. (For modeling data with bounded physical region: $n_{\tilde{g}} \geq 0$, and for application of Bayes' theorem see e.g. Sec. 28 of [21].) Finally, one can also fix the number of gluinos and determine $\alpha_s(M_Z)$. For $n_{\tilde{g}} = 0$ (i.e. QCD) we get $.124 \pm .004$, for $n_{\tilde{g}} = 1$ we get $.129 \pm .006$ (window I) and $.132 \pm .006$ (window III). As expected [21] the QCD value is slightly larger than the world average.




FIG. 3. The 68.3%, 95.4% and 99.7% CL regions of the combined analysis for the theory with window III light gluino ($M_{\tilde{g}} = 3$ GeV) and the 68.3% CL region for the theory without gluino.

Fig. 2 shows the β -function coefficients with 68.3% and 95.4% CL regions for models without gluinos at M_Z . These curves represent merely a transformation of the curves in the (x, y) plane, they result in similar conclusions for QCD. As it can be seen the value of β_0 is quite well constrained; however, not even the sign of β_1 could be predicted. This result reflects the fact that in the energy region studied the running of the coupling is determined almost exclusively by the one-loop term, proportional to β_0 . An interesting possibility is to fix the cross sections, β_1 , β_2 and the flavour dependence of β_0 to their perturbative QCD values and extract β_0 from the fits. The result is $\beta_0 = 5.70 \pm .64$ to be compared with the theoretical value of 5.75. Following [2] we fixed β_0 and β_1 to their theoretical (QCD) value and performed a fit with β_2 as a free parameter. The allowed region for β_2 contains the theoretical prediction, but the error is an order of magnitude larger than the theoretical value.

Comparing our Fig. 1 and Fig. 10 of [1] –taking proper care of using the same variables– one observes that the overlap of the 68.3% CL regions for the two analyses is quite small. This opens the possibility for a much stronger restriction on the theories without and with gluino. We have parametrized the χ^2 of the multi-jet analysis as a function of x and y , and performed a new fit adding it to our χ^2 function. In the actual analysis we have included the new results in [26]. Since the 4-jet analysis is based on a tree-level calculation we have included the unknown higher order QCD and mass effects. We assumed much larger uncalculated higher order corrections than used in the experimental papers. Due to them we enlarged the axes of the error ellipse by 12% of the theoretical x and y values (relative correction of $\mathcal{O}(\alpha_s)$.) For the 4-jet analyses we have used the mass effects calculated by [25]. The results of the combined analysis for QCD and window III gluinos are presented in Fig. 3. Our result is consistent with QCD + no gluino scenario, while the theory with window III light gluino is excluded on the 99.99 (99.89)% CL for $M_{\tilde{g}} = 3$ (5)

GeV. The window I gluino is excluded on the 99.97% CL.

Again one can fix the underlying gauge group to $SU(3)$ and consider the number of gluinos as a parameter to be fitted. We get $n_{\tilde{g}} = -.35 \pm .33$ for window I and $n_{\tilde{g}} = -.156 \pm .27$ ($n_{\tilde{g}} = -.197 \pm .32$) for window III $M_{\tilde{g}} = 3$ (5) GeV gluinos. Using Bayes' theorem assuming that $n_{\tilde{g}}$ is non-negative integer number, the exclusion CL's are 99.96% for window I and 99.96 (99.76)% for window III gluinos.

Leaving out R_τ from the window III combined analyses, the CL values change only negligibly.

In summary, we have presented a method to analyse the running of α_s from M_τ to M_Z . We have determined the group coefficients of the theory, which are in agreement with QCD. The β -function coefficients are also given with a similar result. Furthermore, our results rule out the QCD+light gluino scenario on the 70.8% and 90.7–93.0% CL for window I and III, respectively. Combining the analysis of the present letter and the method using multi-jet data (cf. ref. [1,26], however treating the theoretical errors more conservatively than the experimental groups) we have got an order of magnitude smaller CL regions for the allowed group coefficients, which supports QCD within 1σ and excludes the QCD+light gluino scenario with a 99.97% for window I and at least 99.89% CL for window III gluinos. Performing a one parameter fit to determine the number of gluinos, we get $n_{\tilde{g}} = -.35 \pm .33$ and $n_{\tilde{g}} = -.197 \pm .32$ for window I and III, respectively. In terms of CL's this corresponds to 99.96% and 99.76%. Needless to say, our results are valid assuming the light gluino extension of the standard model. We have not investigated whether additional new physics could change our exclusion CL limits.

The details of the above analysis, the convergence features and the statistical analysis will be presented in a forthcoming publication [27].

We thank M. Drees, B. Gary, A.L. Kataev and R. Sommer for discussions. This work was partially supported by Hung. Sci. grants OTKA-T16248,T22929.

-
- * On leave from Institute for Theoretical Physics, Eötvös University, H-1088 Budapest, Hungary
- [1] M. Schmelling, in Proc. of 15th Int. Conf. Physics in Collision, Ed. Frontieres, Gif-Sur-Yvette, p. 287 (1995).
 - [2] M. Girone, M. Neubert, Phys. Rev. Lett. **76**, 3061 (1996).
 - [3] G. Farrar, P.Fayet, Phys. Lett. **B76**, 575 (1978).
 - [4] G. Farrar, Phys. Rev. **D51**, 3904 (1995).
 - [5] C. Clavelli, P.W. Coulter, Phys. Rev. **D51**, 1117 (1995).
 - [6] M.B. Cakir, G. Farrar, Phys. Rev. **D50**, 3268 (1994).
 - [7] I. Antoniadis, J. Ellis, D.V. Nanopoulos, Phys. Lett. **B262**, 109 (1991).
 - [8] M. Jezabek, J.H. Kühn, Phys. Lett. **B301**, 121 (1993).
 - [9] J. Ellis, D.V. Nanopoulos, D.A. Ross, Phys. Lett. **B305**, 375 (1993); M. Schmelling, R.D. St.Denis, Phys. Lett. **B329**, 393 (1993).
 - [10] B.A. Campbell, J. Ellis, S. Rudaz, Nucl. Phys. **B198** 1 (1982).
 - [11] G. Altarelli, P. Nason, G. Ridolfi, Z. Phys. **C68**, 257 (1995).
 - [12] M. Neubert, Nucl. Phys. **B463**, 499 (1996).
 - [13] M. Shifman, Int. J. Mod. Phys. **A11**, 3195 (1996).
 - [14] S.G. Gorishni, A.L. Kataev, S.A. Larin, Phys. Lett. **B259**, 144 (1991); L.R. Surguladze, M.A. Samuel, Phys. Rev. Lett. **66**, 560 (1991), erratum ibid. **66**, 2416 (1991).
 - [15] L. Clavelli, P.W. Coulter, L.R. Surguladze, Phys. Rev. **D55**, 4268 (1997).
 - [16] W. Bernreuter, W. Wetzel, Nucl. Phys. **B197**, 228 (1982).
 - [17] L.J. Clavelli, L.R. Surguladze, Phys. Rev. Lett. **78**, 1632 (1997); K.G. Chetyrkin, Phys. Lett. **B391**, 402 (1997).
 - [18] A. Pich, Nucl. Phys. (Proc. Suppl.) **B39**, 326 (1995).
 - [19] L. Duflot, Nucl. Phys. (Proc. Suppl.) **B40**, 37 (1995); T. Coan et al., Phys. Lett. **B356**, 580 (1995).
 - [20] LEP Electroweak Working Group, LEPEWWG 96-02.
 - [21] Particle Data Group (R.M. Barnett et al.), Phys. Rev. **D54**, 1 (1996).
 - [22] H.J. Behrend et al., Phys. Lett **B183**, 400 (1987).
 - [23] G. D'Agostini, W. de Boer, G. Grindhammer, Phys. Lett. **B229**, 160 (1989); D. Haidt, in "Directions in High Energy Physics, vol. 14.", ed. P. Langacker, World. Sci. Singapore, p. 203 (1995); S. Eidelman, F. Jegerlehner, Z. Phys. **C67**, 585 (1995).
 - [24] A. de Gouvea, H. Murayama, hep-ph/9606449.
 - [25] R. Munoz-Tapia, W.J. Stirling, Phys. Rev. **D49**, 3763 (1994).
 - [26] R. Barate et al., CERN-PPE-97-002.
 - [27] F. Csikor, Z. Fodor, to be published